

Dominant Role for Signal Transduction in the Transcriptional Memory of Yeast *GAL* Genes[∇]

Sharmistha Kundu† and Craig L. Peterson*

Interdisciplinary Graduate Program, Program in Molecular Medicine, University of Massachusetts Medical School, Worcester, Massachusetts

Received 29 December 2009/Returned for modification 28 January 2010/Accepted 25 February 2010

Several recent studies have shown that the transcriptional induction of yeast *GAL* genes occurs with faster kinetics if the gene has been previously expressed. Depending on the experimental regimen, this transcriptional “memory” phenomenon can persist for 1 to 2 cell divisions in the absence of an inducer (short-term memory) or for >6 cell divisions (long-term memory). Long-term memory requires the *GAL1* gene, suggesting that memory involves the cytoplasmic inheritance of high levels of Gal1 that are expressed in the initial round of expression. In contrast, short-term memory requires the SWI/SNF chromatin-remodeling enzyme, and thus, it may involve the inheritance of distinct chromatin states. Here we have reevaluated the roles of SWI/SNF, the histone variant H2A.Z, and components of the nuclear pore in both the short-term and long-term memory of *GAL* genes. Our results suggest that the propagation of novel chromatin structures does not contribute to the transcriptional memory of *GAL* genes, but rather, memory of the previous transcription state is controlled primarily by the inheritance of the Gal3p and Gal1p signaling factors.

The establishment and maintenance of transcriptional states that are heritable to progeny play a central role during the development of multicellular organisms. In many cases a transcriptional state is propagated in the absence of the original inducing signal, suggesting some type of transcriptional “memory.” Likewise, unicellular organisms must rapidly adapt to signals from their microenvironment, and this process often involves the activation of complex transcriptional networks. The ability to pass on a “memory” of altered environmental conditions and, thus, a memory of altered transcription states may provide progeny with a selective advantage. Since chromatin structure plays a key role in determining the on/off state of eukaryotic genes, the inheritance of altered chromatin structures may provide one mechanism for transcriptional memory.

The transcriptional regulation of the *GAL* gene cluster of budding yeast serves as a paradigm for a complex gene regulatory network that also exhibits the phenomenon of transcriptional memory. *GAL* genes can be separated broadly into two groups: the structural genes (*GAL1*, *GAL5*, *GAL7*, and *GAL10*) that encode enzymes to metabolize galactose and regulatory genes (*GAL2*, *GAL3*, *GAL4*, and *GAL80*) that encode products that transport galactose and control the expression of the structural genes. The transcription of many of the *GAL* genes is tightly controlled by the sugar present in the medium: the expressions of *GAL1*, *GAL3*, *GAL4*, *GAL7*, and *GAL10* are repressed by glucose, and most *GAL* genes are induced 3- to 1,000-fold in the presence of galactose (*GAL5* is constitutive) (3, 12, 18, 19, 28).

The galactose-dependent transcriptional activation of *GAL*

genes involves a complex regulatory network. The Gal4 activator binds to one or more sites upstream of each inducible *GAL* gene, but in the absence of galactose, the activation domain of Gal4 is inactivated by an interaction with the Gal80 repressor. When galactose is added to cells, the Gal2 permease imports galactose, and the binding of galactose to the cytoplasmic Gal3 protein allows Gal3 to bind to Gal80, sequestering the repressor in the cytoplasm (3, 17, 28). Notably, the product of *GAL1*, galactokinase, can substitute for Gal3 when present at high concentrations (14, 15, 29). Thus, Gal3 and Gal1 function as key signal transducers that activate Gal4. The Gal2 and Gal3 proteins are expressed at low basal levels in the absence of galactose, and their expression is increased in galactose medium, creating two positive-feedback loops. The 1,000-fold increase in Gal1 expression is likely to further enhance the Gal3 feedback loop. These positive loops are antagonized by a negative-feedback loop involving the Gal80 repressor, which is induced 2- to 3-fold by galactose. The Gal3 and Gal80 loops work in concert to ensure *GAL* gene homeostasis (2, 30).

Several studies have demonstrated that the Gal3/Gal1 feedback loop provides cells with a “memory” of previous galactose exposure (2, 37). For instance, when naïve, glucose-grown cells are switched to galactose medium, the full induction of *GAL* gene transcription is rather slow, requiring 2 to 4 h (21). These slow induction kinetics presumably reflect the need to synthesize additional Gal4 and to overcome glucose repression mechanisms that occur in *cis* at *GAL* genes (19). However, if cells were previously exposed to galactose, the reinduction of *GAL* genes that follows a 12-h period of glucose repression occurs with much more rapid kinetics. These rapid reinduction kinetics require the Gal1 protein, and heterokaryon studies indicate that the cytoplasmic inheritance of Gal1 provides the memory of previous galactose exposure (37). Interestingly, Brickner and colleagues also reported that the histone variant H2A.Z (also known as Htz1) is also required for transcriptional mem-

* Corresponding author. Mailing address: Program in Molecular Medicine, University of Massachusetts Medical School, 373 Plantation Street, Worcester, MA 01605. Phone: (508) 856-5858. Fax: (508) 856-5011. E-mail: craig.peterson@umassmed.edu.

† Present address: HHMI/Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA.

[∇] Published ahead of print on 8 March 2010.

ory and, furthermore, that memory correlates with the association of a *GAL* gene with the nuclear periphery (4).

Previously, we also described a transcriptional memory phenomenon at yeast *GAL* genes. In our experimental regimen, cells were grown in a neutral, nonrepressing sugar (raffinose) prior to the first round of transcriptional induction in galactose. In this case, *GAL* transcription reaches maximal levels by ~1 h. *GAL* genes were then repressed by growth in glucose medium for 30 min to 4 h, and after transfer to galactose medium, the reinduction of *GAL* genes was found to reach maximal levels by ~5 min. Recently, Laine et al. used a similar protocol to monitor *GAL* transcriptional memory (22). Those studies indicated that the rapid reactivation of *GAL* genes requires the SWI/SNF chromatin-remodeling enzyme and an intragene loop between the 5' and 3' ends of a *GAL* gene (21, 22). Interestingly, the memory of previous galactose exposure persisted through at least one cell division but was lost by 6 to 8 h of growth in glucose. This result contrasts with data from the studies described above, where memory was stable for at least 12 h (4, 37). Given the key role of the SWI/SNF remodeling enzyme in transcriptional memory, we proposed that SWI/SNF may establish an "active" chromatin structure at *GAL* genes that could persist through a few generations in the absence of the galactose inducer. The roles of H2A.Z, the nuclear periphery, or the Gal1 or Gal3 protein in this "short-term" memory phenomenon have not been tested.

Here we reevaluate the role of chromatin-remodeling factors, the histone variant H2A.Z, the nuclear periphery, and cytoplasmic signaling molecules in both the "short-term" and "long-term" transcriptional memory of *GAL* genes. We find that neither H2A.Z nor the recruitment of *GAL* genes to the nuclear periphery is required for the "short-term" memory of *GAL* genes, nor does recruitment to the nuclear periphery appear to be required for "long-term" memory. In contrast, our data suggest that Gal1 and Gal3 function redundantly to promote rapid reinduction kinetics, likely through their cytoplasmic inheritance. Furthermore, SWI/SNF does not appear to be involved in memory *per se*, but rather, its chromatin-remodeling activity is required in both rounds of induction to achieve rapid activation. However, the function of SWI/SNF is most apparent during reinduction, when rapid signaling by Gal3/Gal1 renders chromatin remodeling a rate-limiting step for *GAL1* activation.

MATERIALS AND METHODS

Yeast strains, media, and culture conditions. Strains used in this study are isogenic derivatives of the S288c background. *Saccharomyces cerevisiae* liquid cultures were grown at 30°C in YEP (1% yeast extract, 2% Bacto-peptone) medium supplemented with 2% glucose, 2% galactose, or 2% raffinose plus 0.2% sucrose depending on whether *GAL1* activation or repression was required. For Gal3p-overexpressing strains, wild-type and *swi2Δ* strains were transformed with 2- μ m plasmids expressing full-length Gal3p from a constitutive *ADHI* promoter or the relevant vector control. Transformants were selected and grown on synthetic media (0.67 g yeast nitrogen base without amino acids per 100 ml) containing 2% dextrose lacking uracil (URA).

RNA isolation and analysis. Total RNA was isolated from yeast cells grown to logarithmic phase in appropriate media by the hot-phenol extraction method. The concentration of RNA was estimated by measuring the A_{260} after dissolving it in diethyl pyrocarbonate-treated water. Ten micrograms of total RNA from each sample was electrophoresed on 1% formaldehyde agarose gels, and Northern blotting was done. The housekeeping gene *ACT1* was used as a loading control. Radioactively labeled probes for hybridization were generated by PCR

amplifying the complete *GAL1*, *GAL10*, or *ACT1* open reading frame (ORF) from genomic DNA.

RT-PCR. Cells were grown to mid-log phase in YEP medium with 2% glucose, 2% galactose, or 2% raffinose plus 0.2% sucrose at 30°C. Ten milliliters of cells was harvested, and total RNA was extracted as described above. First-strand cDNA was synthesized by using 2.5 μ g RNA, Superscript II RNase H⁻ reverse transcriptase (RT) (Invitrogen), and 2 pmol each downstream primer designed for genes of interest, according to the manufacturer's instructions. Subsequently, ³²P-labeled PCR was performed by using 2 μ l of the first-strand cDNA reaction mixture and gene-specific primer sets to determine the relative levels of *GAL1*, *GAL3*, *GAL10*, and *ACT1* mRNA for each strain. After 12 cycles (for *GAL1*, *GAL10*, and *ACT1*) or 25 cycles (for *GAL3*) of amplification, PCR products were electrophoresed on 10% acrylamide gels. Reactions were visualized by PhosphorImager (Molecular Dynamics) analysis and quantified by use of Image-Quant software (Amersham Biosciences).

ChIP. Rabbit polyclonal antibody to the C terminus of histone H3 (ab1791) was obtained from Abcam, Inc. Mouse monoclonal antibody to RNA polymerase II (RNAPII) (CTD4H8) was obtained from Covance Research Products. Chromatin immunoprecipitation (IP) (ChIP) assays were performed as described previously by Li et al. (25). The immunoprecipitated DNA was amplified by using quantitative PCR performed with [α -³²P]dCTP and then electrophoresed on 5% acrylamide gels. Reactions were visualized and quantified by PhosphorImager analysis.

Mononucleosome preparation. Samples (from 100 ml of culture at an A_{600} of ~0.7) were cross-linked for 30 min with 37% formaldehyde (final concentration of 2%) at 30°C. Reactions were quenched by the addition of 2.5 M glycine to a final concentration of 125 mM. Cell pellets were collected and washed with water to remove residual medium. Mononucleosomes were prepared as described previously by Dion et al. (9). An aliquot of this sample was deproteinized, and cross-links were reversed. Phenol-chloroform extraction was done, and samples were ethanol precipitated. The resulting pellet was resuspended and treated with RNase A (1 μ g for 1 h at 37°C) to remove all RNA. Samples were then electrophoresed on 1.5% agarose gels to determine the best titration that yielded mononucleosomal DNA.

Nucleosome-scanning analysis. The method for nucleosome-scanning analysis was adapted from a method described previously by Sekinger et al. (32), with modifications. Briefly, mononucleosomal chromatin was prepared as described above. This material was used for IP with histone H3 antibody. The immunoprecipitated DNA was amplified by using quantitative PCR and a set of overlapping primer pairs that were staggered 20 bp relative to each other and together covered an approximately 300-bp region of DNA. The products of all primer pairs were approximately 100 bp long. The efficiency of each primer pair was assayed by performing quantitative PCR with genomic DNA.

RESULTS

Short-term memory of *GAL* genes requires Gal3 and Gal1. Transcriptional memory at *GAL1* has been defined as the ability to reinduce *GAL1* transcription with much faster kinetics than induction rates for the initial exposure to galactose. In our previous study, we showed that memory persisted through only a few cell divisions during glucose repression (21). In contrast, two groups described a different experimental regimen in which *GAL* transcriptional memory persisted for at least 12 h in glucose medium (~4 to 6 cell divisions) (4, 37). In these cases, the initial round of *GAL* induction involved the transfer of repressed, glucose-grown cultures to galactose medium, whereas in our case, raffinose-grown cultures were shifted to galactose. Given the apparent differences in the durations of memory between these studies, we tested whether our strains could recapitulate long-term memory.

Single colonies of isogenic strains were isolated from glucose plates and grown overnight in glucose medium. *GAL10* transcription was then induced by switching cultures to galactose medium (Glc→Gal), and cell aliquots were removed at various times for RNA isolation. Following 10 h in galactose medium, cells were transferred to repressive glucose medium for 12 h, followed by the reinduction of *GAL* genes by a switch back to




FIG. 1. Long-term memory at *GAL* genes requires Gal1p but not SWI/SNF. The schematic at the top depicts a regimen of growth in different carbon sources. Gal, 2% galactose; Glc, 2% glucose. (a) RT-PCR analysis of *GAL10* mRNA levels following initial induction and reinduction after long-term glucose repression (12 h). The wild-type (WT) culture showed a memory of previous *GAL10* induction, which was lost in the *gal1Δ* strain. (b) RT-PCR of *GAL1* and *GAL10* mRNA levels from the *swi2Δ* strain. The reinduction of *GAL1* and *GAL10* is fast compared to initial induction following long-term glucose repression (12 h). Data were averaged over three independent experiments and are represented as relative fold increases over *ACT1* mRNA normalized to a maximum value of 1. Error bars represent standard deviations at each point.

galactose medium (Glc→Gal→Glc→Gal). RT-PCR analysis of *GAL10* transcripts demonstrated that the initial round of expression is quite slow, with peak levels of expression in both the wild-type and *gal1Δ* strains requiring 6 to 7 h (Fig. 1a). In contrast, the reinduction of *GAL10* in the wild-type strain was rapid, with peak levels of expression occurring within 2 to 3 h (Fig. 1a, left). Consistent with previous results, these rapid reinduction kinetics required *GAL1* (Fig. 1a, right). In contrast, the inactivation of the SWI/SNF remodeling complex (*swi2Δ*) showed a rapid reinduction of both *GAL10* and *GAL1* (Fig. 1b and data not shown). Thus, in contrast to our previous study that demonstrated SWI/SNF dependence for the short-term memory of *GAL* gene transcription (21; see also Fig. 6), SWI/SNF is not required for long-term memory. This result is consistent with the cytoplasmic inheritance of Gal1 playing a dominant role in long-term memory (37).

Next, we tested whether Gal1p was also required for short-term memory. In this case, raffinose-grown cells were switched to galactose medium for the initial induction. In this case, maximal expression occurred at between 1 and 2 h. After 2 h, glucose was added to repress *GAL* expression for 60 min, and cells were then switched back to galactose medium (Raf→Gal→Glc→Gal). As expected, wild-type cells showed robust short-term memory at *GAL10*, with maximal expression occurring within 10 min (Fig. 2). Surprisingly, short-term mem-

ory was not eliminated in the *gal1Δ* strain, although the kinetics were reproducibly slower than those of the wild-type strain. Thus, it appears that Gal1p is not essential for the transcriptional memory of *GAL10* expression after a brief period of repression.

During the initial induction with galactose, the *GAL3* gene was induced ~3-fold, whereas *GAL1* was induced ~1,000-fold. Thus, we considered the possibility that memory involves the cytoplasmic inheritance of both Gal3 and Gal1; however, after many generations of growth in glucose (long-term memory), only Gal1 contributes to memory, as the considerably lower level of induced Gal3 would be depleted within a few cell divisions. Consistent with this view, *GAL3* is not required for long-term memory (37). In contrast, during a short-term memory experiment, Gal3 and Gal1 might function in a redundant fashion. To test this possibility, we used strains in which Gal3 is expressed from a doxycycline-inducible promoter (*P_{ter}-GAL3*). In this strain, Gal3 was no longer inducible by galactose, but expression was maintained at a low, basal level in the absence of doxycycline (Fig. 3a) (2). In the absence of doxycycline, sufficient Gal3 is expressed to support a normal induction of *GAL* genes when cells are switched from raffinose to galactose medium (Fig. 3b). Likewise, when cells were then switched to glucose for 60 min, the reinduction of *GAL1* was only slightly slower than that of the wild-type strain, indicating




FIG. 2. Short-term memory at *GAL10* does not require Gal1p. Shown is Northern analysis of *GAL10* RNA levels in the wild-type (left) and *gal1 Δ* (right) strains during initial induction and reinduction following short-term glucose repression (1 h). The *gal1 Δ* strain rapidly reinduces *GAL10* transcription with kinetics almost similar to those of the wild type. The graph shows data from an average of three experiments, while the top shows a representative experiment. Error bars represent standard deviations. *ACT1* is the loading control for total RNA levels.

that memory was partially intact in the absence of Gal3 induction (Fig. 3b). However, the deletion of the *GAL1* gene in the *P_{tet}-GAL3* strain eliminated the rapid reinduction of *GAL10*, indicating that Gal1 and Gal3 play overlapping roles in transcriptional memory (Fig. 3c). In this strain, we also reproducibly observed a decrease in *GAL10* transcription at later time points during the first induction and a poor accumulation of transcripts even at late time points in the second round. This is likely due to the galactose-dependent induction of the Gal80 negative-feedback loop. Since Gal3 levels were not concomitantly induced in these strains, the imbalance in the Gal80/Gal3 ratio is expected from previous studies to prevent Gal4 activation (see references 2 and 30 and references therein). These data further support the view that Gal1 and Gal3 function as redundant inducers of the *GAL* system and that this positive-feedback loop plays a major role in transcriptional memory.

Transcriptional memory of *GAL* genes does not require tethering at the nuclear periphery. Several studies demonstrated that the activation of *GAL1* or *GAL10* transcription leads to the migration of the locus to the nuclear periphery, where it interacts with nuclear pore complexes (NPCs) (1, 4–6, 34). This relocalization event requires several components of the nuclear pore complex, such as Nup2 and Mlp1 (1, 4, 35, 36), as well as nuclear pore-associated factors that regulate mRNA export, such as Sac3 (8, 11, 16, 23). Initially, we tested whether the Nup2-dependent tethering of *GAL10* to the nuclear periphery is required for long-term memory. We monitored *GAL1* induction and reinduction kinetics in a *nup2 Δ* strain (Fig. 4a) and found that the inactivation of Nup2 had no

significant effect on the long-term transcriptional memory of *GAL10*.

We then tested whether relocalization to the nuclear periphery plays a role in a short-term memory regimen. In these experiments, an isogenic set of strains that lacked either Nup2 or Sac3 was grown in galactose and then switched to glucose medium for 1 h prior to reinduction by the addition of galactose. Both of these components were previously shown to play essential roles in the relocalization of active *GAL* genes to the nuclear periphery (5, 20). As shown previously, high levels of *GAL1* transcripts are detectable ~ 10 min after the addition of galactose in the wild-type strain. Strikingly, the short-term memory of *GAL* transcription was intact in each of the mutants that blocked *GAL* localization to the periphery (Fig. 4b and data not shown). Thus, localization to the nuclear periphery is not essential for rapid *GAL1* reinduction. These results are consistent with several studies demonstrating that plasmid-borne *GAL* genes are released from the nuclear periphery following >1 h of growth in glucose medium (1, 36; S. Kundu, C. L. Peterson, and M. Rosbash, unpublished results), and thus, peripheral localization in most strains does not appear to be maintained through more than one cell cycle.

H2A.Z does not contribute to short-term memory. Brickner and colleagues (4) previously reported that the long-term memory of *GAL1* transcription requires the histone variant H2A.Z (also known as Htz1). In our strain background (s288C), we found that an *htz1 Δ* mutant had a significant defect in the initial induction of *GAL1* or *GAL10* when cells were switched from glucose to galactose medium, and this transcriptional defect made analyses of long-term memory problematic




FIG. 3. Cytoplasmic signaling by both Gal1p and Gal3p is required for optimal *GAL* gene reinduction. (a) RT-PCR of wild-type and *gal3Δ;P_{tet}-GAL3* strains showing *GAL3* expression levels in glucose or galactose. (b) Minimal expression of Gal3p from a *P_{tet}-GAL3* construct causes a partial defect in *GAL1* reinduction following short-term glucose repression (1 h). Shown are *GAL1* mRNA levels for the wild-type (left) and *gal3Δ-P_{tet}-GAL3* (right) strains. At top is a representative RT-PCR, and the graphs below show data from an average of three independent experiments represented as fold increases over *ACT1* mRNA normalized to a maximum value of 1. Error bars show standard deviations at each point. o/n, overnight. (c) *GAL10* is induced in minimally expressing *P_{tet}-GAL3* and an amino-terminal deletion of a *GAL1* double mutant but is lost at 2 h postinduction. *GAL10* reinduction is severely compromised even after 8 h following short-term glucose repression (1 h). Data in panel c are representative of five independent experiments.

(data not shown). We found similar results for strains of the w303 background (data not shown). Our observations are consistent with data from a previous study demonstrating that Htz1 plays a key role in the recruitment of the Mediator complex during the initial activation of a glucose-repressed *GAL* gene (13, 24). To circumvent this issue, we asked if Htz1 was required for short-term memory. When cells were switched from raffinose to galactose medium, both the wild-type and *htz1Δ* strains showed identical kinetics of initial *GAL1* induction (Fig. 4C). Interestingly, the *htz1Δ* strain also demonstrated a rapid reinduction of *GAL1* transcription, indicating that Htz1

is not required for short-term memory in strains of the s288C background (Fig. 4C).

SWI/SNF promotes rapid PIC loading but does not generate alternate nucleosome positions. These studies indicate that SWI/SNF and Gal1/Gal3 may be the primary factors that control the transcriptional memory of *GAL* genes. To test if chromatin remodeling by SWI/SNF facilitates faster preinitiation complex (PIC) assembly on the *GAL1* promoter during reinduction, we monitored the RNA polymerase II (RNAPII) association with the *GAL1* promoter during a cycle of expression, repression, and reinduction. For both the wild-type and




FIG. 4. *GAL1* memory does not require nuclear pore localization or H2A.Z. Schematics at the top depict regimens of growth in different carbon sources. Raf, 2% raffinose; Gal, 2% galactose; Glc, 2% glucose. (a) RT-PCR analysis of *GAL10* induction and reinduction in *nup2Δ* cells following long-term glucose repression (12 h). Long-term memory does not require Nup2p. Data represent data from three independent experiments. (b) Short-term *GAL1* memory is unaffected in *nup2Δ* (left) and *sac3Δ* (right) mutants. (c) Northern analysis of wild-type (left) and *htz1Δ* (right) strains showing rapid *GAL1* reinduction following short-term glucose repression (1 h). Data are representative of data from three independent experiments.

swi2Δ strains, RNAPII was rapidly lost from the promoter during glucose repression, as observed previously by us and others (21, 31). Glucose repression is also associated with a rapid reassembly of nucleosomes over the *GAL1* promoter in both wild-type and *swi2Δ* strains (Fig. 5a, left). During reinduction in the wild-type strain, RNAPII was recruited to *GAL1* within 10 min, paralleling the rapid appearance of *GAL1* transcripts (Fig. 5a, right). Likewise, both TATA binding protein (TBP) and SWI/SNF were rapidly recruited to the *GAL1* locus during reinduction, and transcriptional reinduction was associated with a rapid nucleosome loss (data not shown). In contrast, RNAPII was recruited slowly in the *swi2Δ* strain, reflecting the requirement for SWI/SNF to act at an early step in

promoting rapid RNAPII re-recruitment during *GAL* gene reinduction.

In our previous study, we proposed a model in which SWI/SNF might control transcriptional memory by influencing the positioning of nucleosomes that are reassembled onto the *GAL1* promoter during glucose repression. Since SWI/SNF rapidly dissociates from the glucose-repressed *GAL1* promoter (21), this model further proposes that these alternative nucleosome positions are propagated through DNA replication in the absence of SWI/SNF. Changes in nucleosome positioning might then enhance the rate of RNA polymerase II recruitment. To test this model directly, we mapped the positions of two promoter-proximal nucleosomes at the *GAL1* locus (see




FIG. 5. SWI/SNF promotes RNA polymerase II loading but does not generate alternate nucleosome positions at the *GAL1* promoter. (a, left) Histone H3 ChIP of the wild-type and *swi2Δ* strains to measure nucleosome occupancy at the *GAL1* promoter. The loss of SWI/SNF does not inhibit the kinetics of promoter nucleosome reloading during glucose repression. (Right) RNA polymerase II ChIP of the wild-type and *swi2Δ* strains showing that the faster recruitment during *GAL1* reinduction is dependent on SWI/SNF. RNA polymerase II and H3 levels were tested at the *GAL1* promoter and normalized to a telomere sequence (Chr VI, 70 bp from the right end). (b) Schematic representation of the *GAL1-10* regulatory region. UAS_{GAL} marks the Gal4p binding sites. URS_{GAL} is the binding site for the glucose-dependent repressor Mig1p. Ovals represent previously mapped nucleosome positions. Ovals shown as solid lines represent *GAL1* promoter nucleosomes that are mapped in c and d. TATA represents the TBP binding sites, and +1 represents the transcription start sites. (c) Nucleosome-scanning ChIP with histone H3 antibody in the wild-type strain for promoter nucleosomes B (left) and C (right). Black lines represent cultures grown in glucose overnight, and gray lines represent short-term (1-h) glucose-repressed cultures following a brief *GAL1* induction. On the x axes of graphs, B1 to B10 represent 10 primer pairs spanning positions -302 to +3 from the translation start site. C1 to C10 represent nine primer pairs spanning positions -148 to +160 from the translation start site. On the y axis, the relative percent IP of H3 normalized to a maximum value of 1 is plotted. (d) Same as panel c but with an *swi2Δ* strain.

Fig. 5b for a schematic representation) in wild-type and *swi2Δ* cells. Nucleosome positioning was mapped under two conditions: (i) cells that were grown continuously in glucose (long-term repression) and (ii) cells that had been grown in galactose and then repressed with glucose for 1 h (short-term repression). In each case, cells were treated with formaldehyde prior to collection and analysis of nucleosome positioning by nucleosome-scanning ChIP (32). Mononucleosomal chromatin was prepared by MNase digestion and used for ChIP with an anti-histone H3 antibody. Quantitative PCR was performed with primer pairs scanning approximately 300 bp around the predicted dyads of promoter nucleosome B (Fig. 5c, left) and promoter nucleosome C (Fig. 5c, right).

In long-term-repressed cells, nucleosomes B and C were positioned as predicted from previous studies, each protecting ~160 bp of DNA. In contrast, no ChIP signal was detected when mononucleosomal chromatin was prepared from cells growing in galactose, consistent with a loss of *GAL1* promoter nucleosomes (26, 27). However, when galactose-grown cells were exposed to glucose for 1 h, nucleosomes were reassembled, and the positioning of NucB and NucC appeared to be nearly identical between long-term-repressed and short-term-repressed cultures. Notably, this result is consistent with previous analyses of the *GAL1* chromatin structure (7, 26). We repeated the same set of experiments with an *swi2Δ* strain and obtained similar results, confirming that SWI/SNF was not involved in nucleosome positioning at the *GAL1* promoter during glucose repression (Fig. 5d). Thus, these data indicate that the role of SWI/SNF in promoting rapid *GAL1* reinduction does not involve the generation of a novel pattern of nucleosome positions at the *GAL1* promoter.

Gal3p coinducer overexpression leads to rapid *GAL1* induction in wild-type cells. We considered an alternate hypothesis in which SWI/SNF functions in both the first and second rounds of *GAL1* induction but SWI/SNF action is only rate limiting for expression during reinduction. This model proposes that *GAL1* expression is controlled by two different kinetic steps: the accumulation of signaling molecules, Gal1 and Gal3 (designated k_1), which antagonize the Gal80 repressor, and chromatin remodeling (designated k_2) (see model in Fig. 7). In this model, the accumulation of signaling molecules (k_1) is the rate-limiting step for the initial induction of *GAL* genes (i.e., $k_2 > k_1$). However, in the second round of expression, signaling is very rapid due to the high levels of Gal3p and Gal1p that accumulate during initial induction, and chromatin remodeling by SWI/SNF becomes the slow step (i.e., $k_2 < k_1$). Note that in this model, SWI/SNF action must be partially redundant with other remodeling enzymes that remodel the promoter, albeit more slowly, in its absence (see Fig. 7). SWI/SNF is not required for the induced levels of Gal3 or Gal1, eliminating simple models where SWI/SNF directly regulates the Gal1/Gal3 feedback loop (18).

One prediction of this model is that increasing the levels of signaling molecules prior to the first round of induction should uncover a kinetic role for SWI/SNF during the initial round of expression. To test this idea, both wild-type and *swi2Δ* strains were transformed with a high-copy-number plasmid that harbors a constitutively expressed *ADHI-GAL3* gene. The growth of these cells in raffinose medium leads to high levels of *GAL3* expression in both the wild-type and *swi2Δ* strains (Fig. 6a). In

the wild-type strain, the constitutive expression of Gal3p increased the rate of the initial induction such that the accumulation of *GAL1* transcripts was nearly indistinguishable from the rate of reinduction (Fig. 6b, right). In contrast, cells transformed with the vector control showed slow kinetics of *GAL1* expression in the first round and rapid kinetics during reinduction (Fig. 6b, left). Thus, constitutive, high-level expression of Gal3 appears to obviate transcriptional memory. Strikingly, the constitutive expression of *GAL3* uncovered a significant role for SWI/SNF during the first round of expression (Fig. 5c, right). In fact, the *GAL1* induction rates in the *swi2Δ* strain were identical in the first and second rounds of expression and were slow compared to wild-type rates. Thus, SWI/SNF action is generally rate limiting for *GAL* expression when cells contain high levels of Gal3. These data are fully consistent with the kinetic model shown in Fig. 7.

DISCUSSION

In the present study, we have carried out experiments aimed at reconciling studies that have investigated transcriptional memory at yeast *GAL* genes. In our previous work, we showed that the SWI/SNF remodeling enzyme was required for the rapid reinduction of *GAL1* transcription following a 0.5- to 4-h period of glucose repression. In our experimental regimen, transcriptional memory was inherited through only 1 to 2 cell divisions. In contrast, two groups reported an experimental scheme in which the transcriptional memory of *GAL* genes was maintained for at least 12 h under glucose-repressed conditions (4 to 6 divisions). In this case, memory was shown to require a *GAL1* gene product, the H2A.Z histone variant, and it correlated with peripheral nuclear localization. Here we have shown that the tethering of *GAL* loci to the nuclear periphery is not essential for transcriptional memory in either scheme. Our data also indicate that SWI/SNF, and likely H2A.Z, functions downstream of the transcriptional memory process, regulating the general kinetics of *GAL* induction. In contrast, transcriptional memory generally requires the Gal1/Gal3 feedback loop, with the function of Gal1, as a weak Gal3-like inducer, predominating in the long-term memory regimen and both factors showing overlapping functions in a short-term memory scheme. Thus, the transcriptional memory of *GAL* genes does not appear to involve the inheritance of chromatin states but rather involves the cytoplasmic inheritance of Gal1 and/or Gal3 signaling factors.

The kinetics of *GAL* gene induction are strongly influenced by the concentration of galactose in the medium, levels of the Gal3 inducer, and levels of the Gal80 repressor (2, 30). Alterations in the levels of these factors can have a strong impact on *GAL* transcription. Likewise, our data indicate that the levels of Gal3 can also impact the extent to which SWI/SNF is required for the optimal kinetics of *GAL* induction. At low, basal Gal3 concentrations, SWI/SNF action does not appear to be rate limiting for *GAL* expression, as the inactivation of SWI/SNF has only a minor effect on induction kinetics (Fig. 1). However, an increased expression of Gal3 leads to a more extensive *GAL* transcriptional defect in the absence of SWI/SNF. These results provide an explanation for why *swi/snf* mutations in some strain backgrounds lead to strong defects in *GAL* transcription even during an initial round of expression




FIG. 6. Gal3p coinducer overexpression leads to rapid *GAL1* induction in the wild type. (a) *GAL3* mRNA expression compared between growth in raffinose and that in galactose for the indicated strains and shown as fold induction over *ACT1*. (b) Northern analysis using the short-term memory regimen for wild-type cells transformed with the control plasmid (left) and a Gal3p-overexpressing plasmid (*P_{ADHI}-GAL3*) (right). (c) Same as panel b for the *swi2Δ* strain. The constitutive expression of Gal3p rapidly activates *GAL1* transcription even upon initial induction in the wild type. The overexpression of Gal3p uncovers a more prominent role for SWI/SNF during the initial round of induction. All Northern blots were subsequently probed for *ACT1* as a loading control.

(10). Presumably, these strain backgrounds have a higher level of basal Gal3 signaling (or lower levels of the Gal80 repressor). Differences in the potencies of the Gal3 positive-feedback loop among strain backgrounds may also explain why the Htz1 histone variant plays variable roles in the initial induction of *GAL* genes (4, 13).

In the strain background used in our studies (s288C), Htz1 contributes significantly to the initial induction kinetics of *GAL1* when cells are transferred from glucose to galactose medium, and consequently, we were unable to assess its role during the long-term memory regimen. We have also found that Htz1 is required for rapid *GAL* induction kinetics using a second, common strain background (w303) (our unpublished observations). Similar results were reported previously by Gligoris et al. (13). In contrast, Brickner and colleagues (4) used the JBY strain background, and thus, it remains a possibility

that Htz1 may contribute to the transcriptional memory phenomenon in these strains.

Several studies demonstrated that the activation of *GAL* gene expression leads to the rapid localization of these loci to the nuclear periphery and interaction with NPCs. It was suggested that this NPC association may facilitate the rapid export of *GAL* mRNA into the cytoplasm. Brickner and colleagues (4) also reported that *GAL1* maintains its association with the NPC for many cell divisions, suggesting that peripheral localization may be involved in memory. The genetic requirements for *GAL1* localization to the nuclear periphery are well established, with key roles being played by the NPC component Nup2 and components of the mRNA processing and export machinery, such as Sac3. We find that the inactivation of any one of these factors has no detectable impact on *GAL* transcriptional memory, indicating that while *GAL* gene transcrip-




FIG. 7. Two-step model for optimum induction kinetics of *GAL1* transcription. Extracellular galactose is transported into the cytoplasm by Gal2p, and galactose-bound Gal3p or Gal1p antagonizes the Gal80 repressor, leading to the activation of the Gal4p activator and allowing it to recruit SWI/SNF. Chromatin remodeling by SWI/SNF leads to the rapid loss of promoter nucleosomes and PIC assembly, resulting in *GAL1* induction. Gal4p activation (shown as Gal4p*) leads to an increased production of Gal1p and Gal3p, thereby amplifying the signal. k_1 and k_2 represent the rate-determining steps of this pathway, where k_1 reflects the action of Gal1p and Gal3p and k_2 reflects chromatin remodeling by SWI/SNF. In the absence of SWI/SNF, other nonspecific chromatin remodelers may cause promoter nucleosome loss. In the initial round of induction, Gal1p and Gal3p levels are low, making k_1 the rate-determining step. After the initial round of expression and a brief period of glucose repression, Gal3p and Gal1p levels are much higher than those in the initial state. Hence, k_1 is fast during *GAL* reinduction, making k_2 the slowest and rate-determining step.

tion tethers the loci to the nuclear periphery, tethering itself is not required for memory. We note that our results are consistent with several studies from the Rosbash group demonstrating that *GAL* genes are released from the periphery after 40 min to 1 h in glucose medium (see, e.g., reference 36) and thus, tethering does not generally correlate well with memory. Likewise, we have used a plasmid-based system in our strain background for the monitoring of *GAL* localization, and we also find that plasmid-borne *GAL* genes are released from the periphery after 1 h of glucose repression (36; S. Kundu, C. L. Peterson, M. Rosbash, and S. Vodala, unpublished results). Furthermore, previous studies have shown that parental and newly assembled NPCs are asymmetrically segregated during yeast cell division. Daughter cells receive only new nuclear membrane material and newly assembled NPCs, with all of the parental NPCs remaining in the mother cell (33). Thus, if a gene locus is associated with a parental NPC, a daughter cell

can inherit this structure from the mother only if interactions with parental and newly assembled NPCs are dynamic. Such dynamic interactions, however, would appear to be inconsistent with a memory phenomenon that requires the faithful segregation of the parental state. Together, the data indicate that the tethering of loci to NPCs is unlikely to contribute to transcriptional memory events that are heritable to progeny.

Recently, two groups reported that the transcriptional induction of *GAL* genes leads to the formation of an intragene loop between the 5' and 3' ends of the *GAL1,10* gene cluster (22, 35). Loop formation requires the general transcription factor TFIIB and components of the mRNA processing and export machinery. During a period of glucose repression, the stability of this gene loop also requires the NPC-associated protein Mlp1. Interestingly, this loop is maintained for at least 1 cell division in glucose medium, and the disruption of the intragene loop leads to the loss of the short-term transcriptional memory of *GAL* genes. SWI/SNF is not required for gene loop formation or maintenance, suggesting that the SWI/SNF action functions downstream. Those authors also showed that the intragene loop is required for rapid RNAPII recruitment during *GAL1* reinduction, leading those authors to suggest that the intragene loop facilitates the reassociation of RNAPII during the second round of expression. However, since SWI/SNF is also required for RNAPII recruitment, but not loop formation, these data indicate that the loop is not sufficient for RNAPII recruitment. The intragene loop is required to maintain the Gal4 activator at the promoter during glucose repression, so one simple model that we favor is that the loop functions primarily to maintain Gal4 so that it is poised to rapidly re-recruit SWI/SNF and other key targets when galactose is reencountered in the environment.

REFERENCES

1. Abruzzi, K. C., D. A. Belostotsky, J. A. Chekanova, K. Dower, and M. Rosbash. 2006. 3'-end formation signals modulate the association of genes with the nuclear periphery as well as mRNA dot formation. *EMBO J.* **25**: 4253–4262.
2. Acar, M., A. Becskei, and A. van Oudenaarden. 2005. Enhancement of cellular memory by reducing stochastic transitions. *Nature* **435**:228–232.
3. Bhat, P. J., and T. V. Murthy. 2001. Transcriptional control of the *GAL1*/MEL regulon of yeast *Saccharomyces cerevisiae*: mechanism of galactose-mediated signal transduction. *Mol. Microbiol.* **40**:1059–1066.
4. Brickner, D. G., I. Cajigas, Y. Fondufe-Mittendorf, S. Ahmed, P. C. Lee, J. Widom, and J. H. Brickner. 2007. H2A.Z-mediated localization of genes at the nuclear periphery confers epigenetic memory of previous transcriptional state. *PLoS Biol.* **5**:e81.
5. Cabal, G. G., A. Genovesio, S. Rodriguez-Navarro, C. Zimmer, O. Gadal, A. Lesne, H. Buc, F. Feuerbach-Fournier, J. C. Olivo-Marin, E. C. Hurt, and U. Nehrbass. 2006. SAGA interacting factors confine sub-diffusion of transcribed genes to the nuclear envelope. *Nature* **441**:770–773.
6. Casolari, J. M., C. R. Brown, S. Komili, J. West, H. Hieronymus, and P. A. Silver. 2004. Genome-wide localization of the nuclear transport machinery couples transcriptional status and nuclear organization. *Cell* **117**:427–439.
7. Cavalli, G., and F. Thoma. 1993. Chromatin transitions during activation and repression of galactose-regulated genes in yeast. *EMBO J.* **12**:4603–4613.
8. Chekanova, J. A., K. C. Abruzzi, M. Rosbash, and D. A. Belostotsky. 2008. Sus1, Sac3, and Thp1 mediate post-transcriptional tethering of active genes to the nuclear rim as well as to non-nascent mRNP. *RNA* **14**:66–77.
9. Dion, M. F., T. Kaplan, M. Kim, S. Buratowski, N. Friedman, and O. J. Rando. 2007. Dynamics of replication-independent histone turnover in budding yeast. *Science* **315**:1405–1408.
10. Ferreira, M. E., P. Prochasson, K. D. Berndt, J. L. Workman, and A. P. Wright. 2009. Activator-binding domains of the SWI/SNF chromatin remodeling complex characterized in vitro are required for its recruitment to promoters in vivo. *FEBS J.* **276**:2557–2565.
11. Fischer, T., K. Strasser, A. Racz, S. Rodriguez-Navarro, M. Oppizzi, P. Ihrig, J. Lechner, and E. Hurt. 2002. The mRNA export machinery requires the novel Sac3p-Thp1p complex to dock at the nucleoplasmic entrance of the nuclear pores. *EMBO J.* **21**:5843–5852.

12. **Frolova, E., M. Johnston, and J. Majors.** 1999. Binding of the glucose-dependent Mig1p repressor to the GAL1 and GAL4 promoters in vivo: regulation by glucose and chromatin structure. *Nucleic Acids Res.* **27**:1350–1358.
13. **Gligoris, T., G. Thireos, and D. Tzamarias.** 2007. The Tup1 corepressor directs Htz1 deposition at a specific promoter nucleosome marking the GAL1 gene for rapid activation. *Mol. Cell. Biol.* **27**:4198–4205.
14. **Hawkins, K. M., and C. D. Smolke.** 2006. The regulatory roles of the galactose permease and kinase in the induction response of the GAL network in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **281**:13485–13492.
15. **Hittinger, C. T., and S. B. Carroll.** 2007. Gene duplication and the adaptive evolution of a classic genetic switch. *Nature* **449**:677–681.
16. **Jani, D., S. Lutz, N. J. Marshall, T. Fischer, A. Kohler, A. M. Ellisdon, E. Hurt, and M. Stewart.** 2009. Sus1, Cdc31, and the Sac3 CID region form a conserved interaction platform that promotes nuclear pore association and mRNA export. *Mol. Cell* **33**:727–737.
17. **Jiang, F., B. R. Frey, M. L. Evans, J. C. Friel, and J. E. Hopper.** 2009. Gene activation by dissociation of an inhibitor from a transcriptional activation domain. *Mol. Cell. Biol.* **29**:5604–5610.
18. **Johnston, M.** 1987. A model fungal gene regulatory mechanism: the GAL genes of *Saccharomyces cerevisiae*. *Microbiol. Rev.* **51**:458–476.
19. **Johnston, M., J. S. Flick, and T. Pexton.** 1994. Multiple mechanisms provide rapid and stringent glucose repression of GAL gene expression in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **14**:3834–3841.
20. **Kohler, A., M. Schneider, G. G. Cabal, U. Nehrass, and E. Hurt.** 2008. Yeast ataxin-7 links histone ubiquitination with gene gating and mRNA export. *Nat. Cell Biol.* **10**:707–715.
21. **Kundu, S., P. J. Horn, and C. L. Peterson.** 2007. SWI/SNF is required for transcriptional memory at the yeast GAL gene cluster. *Genes Dev.* **21**:997–1004.
22. **Laine, J. P., B. N. Singh, S. Krishnamurthy, and M. Hampsey.** 2009. A physiological role for gene loops in yeast. *Genes Dev.* **23**:2604–2609.
23. **Lei, E. P., C. A. Stern, B. Fahrenkrog, H. Krebber, T. I. Moy, U. Aebi, and P. A. Silver.** 2003. Sac3 is an mRNA export factor that localizes to cytoplasmic fibrils of nuclear pore complex. *Mol. Biol. Cell* **14**:836–847.
24. **Lemieux, K., M. Laroche, and L. Gaudreau.** 2008. Variant histone H2A.Z, but not the HMG proteins Nhp6a/b, is essential for the recruitment of Swi/Snf, Mediator, and SAGA to the yeast GAL1 UAS(G). *Biochem. Biophys. Res. Commun.* **369**:1103–1107.
25. **Li, X. Y., S. R. Bhaumik, and M. R. Green.** 2000. Distinct classes of yeast promoters revealed by differential TAF recruitment. *Science* **288**:1242–1244.
26. **Lohr, D.** 1984. Organization of the GAL1-GAL10 intergenic control region chromatin. *Nucleic Acids Res.* **12**:8457–8474.
27. **Lohr, D., and J. Lopez.** 1995. GAL4/GAL80-dependent nucleosome disruption/deposition on the upstream regions of the yeast GAL1-10 and GAL80 genes. *J. Biol. Chem.* **270**:27671–27678.
28. **Lohr, D., P. Venkov, and J. Zlatanova.** 1995. Transcriptional regulation in the yeast GAL gene family: a complex genetic network. *FASEB J.* **9**:777–787.
29. **Meyer, J., A. Walker-Jonah, and C. P. Hollenberg.** 1991. Galactokinase encoded by GAL1 is a bifunctional protein required for induction of the GAL genes in *Kluyveromyces lactis* and is able to suppress the gal3 phenotype in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **11**:5454–5461.
30. **Ramsey, S. A., J. J. Smith, D. Orrell, M. Marelli, T. W. Petersen, P. de Atauri, H. Bolouri, and J. D. Aitchison.** 2006. Dual feedback loops in the GAL regulon suppress cellular heterogeneity in yeast. *Nat. Genet.* **38**:1082–1087.
31. **Schwabish, M. A., and K. Struhl.** 2007. The Swi/Snf complex is important for histone eviction during transcriptional activation and RNA polymerase II elongation in vivo. *Mol. Cell. Biol.* **27**:6987–6995.
32. **Sekinger, E. A., Z. Moqtaderi, and K. Struhl.** 2005. Intrinsic histone-DNA interactions and low nucleosome density are important for preferential accessibility of promoter regions in yeast. *Mol. Cell* **18**:735–748.
33. **Shcheprova, Z., S. Baldi, S. B. Frei, G. Gonnet, and Y. Barral.** 2008. A mechanism for asymmetric segregation of age during yeast budding. *Nature* **454**:728–734.
34. **Taddei, A., G. Van Houwe, F. Hediger, V. Kalck, F. Cubizolles, H. Schober, and S. M. Gasser.** 2006. Nuclear pore association confers optimal expression levels for an inducible yeast gene. *Nature* **441**:774–778.
35. **Tan-Wong, S. M., H. D. Wijayatilake, and N. J. Proudfoot.** 2009. Gene loops function to maintain transcriptional memory through interaction with the nuclear pore complex. *Genes Dev.* **23**:2610–2624.
36. **Vodala, S., K. C. Abruzzi, and M. Rosbash.** 2008. The nuclear exosome and adenylation regulate posttranscriptional tethering of yeast GAL genes to the nuclear periphery. *Mol. Cell* **31**:104–113.
37. **Zacharioudakis, I., T. Gligoris, and D. Tzamarias.** 2007. A yeast catabolic enzyme controls transcriptional memory. *Curr. Biol.* **17**:2041–2046.